

COMMUNICATION

PdBr₂-catalyzed acetal formation of carbonyl compounds using diazophenanthrenequinone: utility of 9,10-phenanthrenedioxy acetal

Mitsuru Kitamura,^{[a]*} Ryo Fujimura,^[a] Tomoaki Nishimura,^[a] Shuhei Takahashi,^[a] Hirokazu Shimooka,^[a] Tatsuo Okauchi^[a]

[a] Prof. Dr. M. Kitamura, R. Fujimura, T. Nishimura, Dr. S. Takahashi, Dr. H. Shimooka, Prof. Dr. T. Okauchi.
Department Applied Chemistry
Kyushu Institute of Technology
1-1 Sensuicho, Tobata, Kitakyushu, 804-8550
E-mail: kita@che.kyutech.ac.jp
Homepage: <http://www.che.kyutech.ac.jp/chem27/chem27.html>

Supporting information for this article is given via a link at the end of the document.

Abstract: We developed a new acetalization method of ketones and aldehydes under non-acidic conditions using diazophenanthrenequinone and PdBr₂. The formed acetals that have a phenanthrene skeleton withstand under mild acidic conditions. Removal of acetals was successfully proceeded under strong acidic or oxidation conditions using aqueous ceric ammonium nitrate (CAN) to afford corresponding ketones and aldehydes.


For the synthesis of complex and functional organic compounds, the carbonyl group is one of the useful functional groups. They are used as both nucleophiles and electrophiles, and various carbon-carbon bond formation reactions using carbonyl compounds are reported.^[1] For example, carbonyl groups suffer nucleophilic attacks at carbon atoms of carbonyl groups owing to carbon nucleophiles such as organometallic reagents and enolates. In contrast, enolates and enols formed by deprotonation and tautomerization from corresponding carbonyl compounds are used as carbon nucleophiles that react with carbon electrophiles such as alkyl halides and carbonyl compounds. However, such high reactivity of carbonyl compounds sometimes interrupts the desired reaction in the multistep synthesis of organic compounds. Side reaction of a carbonyl group can be inhibited *via* protection/deprotection processes in the group, and acetal is usually used to protect the carbonyl group.^[2]

Acetalization of carbonyl compounds is generally conducted *via* dehydration reaction of alcohol under acidic conditions.^[2] However, when the carbonyl compounds have an acid-sensitive functional group, acidic acetalization is difficult to apply to the molecules. Therefore, various methods are being introduced in developing non-acidic acetalization,^[3-6] and several practical methods have been developed, such as iodine promoted reaction,^[3] base-catalyzed reaction,^[4] and metal-catalyzed reaction.^[5]

We have studied the reaction of metal carbene formed from diazoquinones.^[7,8] We envisioned that a practical acetal formation could be developed *via* the reaction between diazoquinones and carbonyl compounds in the presence of a metal catalyst and therefore experimented. In this letter, we describe the outcome of the abovementioned investigation.

In Scheme 1, the outline of our strategy toward acetalization of carbonyl compounds under non-acidic reaction conditions is



depicted and conducted using diazoquinone **1** and a metal catalyst. Diazoquinone **1** reacts with metal catalysts to form metal carbene **I**,^[9] that is attacked with a carbonyl compound **2** to form oxonium ylide **II**.^[10] Internal cyclization proceeds to afford the corresponding acetal **3**. As the diazoquinone for the acetalization, diazophenanthrenequinone **1** was chosen because two outside benzene rings in phenanthrene are responsible for the cyclization to force the reaction sites close, as shown in **II** (vs **II'**). Moreover, the acetal carbon in acetal **3** is not a chiral center, which helps in the multistep transformation so that it does not lead to the formation of a diastereomer *via* the protection step (acetalization).



Scheme 1. Plan of metal-catalyzed acetalization using diazophenanthrenequinone **1**.

Diazophenanthrenequinone **1** was prepared by diazo-transfer reaction of 9-phenanthrol **4** with 2-azido-1,3-dimethylimidazolium chloride (ADMC **5**) in good yield (Scheme 2).^[11]

COMMUNICATION

**Scheme 2.** Preparation of diazophenanthrenequinone **1**.**Table 1.** Optimization study of metal-catalyzed acetalization using diazophenanthrenequinone **1**.^[a]


Run	MLn (x mol%) ^[b]	yield (%) ^[c]	
		3a	6a
1	PdBr ₂ (10)	68	10
2	PdCl ₂ (10)	55	5
3	PdI ₂ (10)	50	12
4	Pd(acac) ₂ (10)	16	trace
5	Pd(OAc) ₂	26	5
6	Pd ₂ (dba) ₃ (10)	23	18
7	Pd ₂ (dba) ₃ (10), PPh ₃ (40)	0	0
8	Rh ₂ (OAc) ₄ (3)	trace	0
9	Rh ₂ (oct) ₄ (3)	6	0
10	Rh ₂ (esp) ₂ (3)	34	trace
11	Cu(CF ₃ -acac) ₂ ·H ₂ O	trace	0

[a] Reaction conditions: **2a** (0.3 mmol), **1** (0.45 mmol), MLn (3–10 mol%) in benzene (1.5 mL) at reflux for 1.5 h. [b] dba = dibenzylideneacetone, acac = acetoacetonato, oct = octanoate, esp = $\alpha,\alpha,\alpha',\alpha'$ -tetramethyl-1,3-benzenedipropionate, CF₃-acac = hexafluoroacetylacetonato. [c] Isolated yield.

An initial trial of the metal-catalyzed acetalization using diazophenanthrenequinone **1** was conducted with 4-phenyl-2-butanone (**2a**) (Table 1). The desired acetalization proceeded efficiently with PdBr₂ (Run 1). When diazophenanthrenequinone **1** (1.5 equiv) in benzene was slowly added to a mixture of ketone **2a** and 10 mol% of PdBr₂ in benzene at reflux, acetalization


proceeded smoothly to afford acetal **3a** in 68% yield. Other palladium(II) halide and palladium(II) salts were ineffective for acetalization compared with PdBr₂ (Runs 1–5). In the reaction with tris(dibenzylideneacetone)dipalladium, acetal **3a** was formed (Run 6), and the formation of **3a** was well suppressed in the presence PPh₃ (Run 7). Although several Rh catalysts were examined, remarkable increase in the yield of acetal **3a** was not observed (Runs 8–10). Cyclization of carbonyl compounds and 2-diazo-1,3-dicarbonyl compounds using Cu(II) salt has been reported^{5b}; however the Cu salt was not effective for the cyclization using diazophenanthrenequinone **1** (Run 11). Additionally, trioxocin derivative **6a** was formed as a by-product in some cases.

The efficiency of PdBr₂ may be attributed to the role of Br⁻. Since 5-*endo-trig* cyclization is highly disfavored commonly,^[12] acetal **3a** is not formed directly from oxonium ylide **IIa**, but is formed by 5-*exo-tet* cyclization of **IIIa** which was formed by the reaction of **IIa** and Br⁻ (Scheme 3).


**Scheme 3.** Efficiency of Br⁻ for the cyclization.

Next, a variety of aldehydes and ketones were examined for the PdBr₂-catalyzed acetalization using diazophenanthrenequinone **1** (Table 2). Both the alkyl and aryl aldehydes were transformed to corresponding acetals in good to high yields. In the series of ketones, cyclic ketones were effective for the acetalization than acyclic ketones in most cases. It is important to note that acid-sensitive TBS group is tolerated under the reaction conditions as shown in the formation of **3h**.

Subsequently, experiment on stability under various conditions and deprotection of acetal **3** was performed (Table 3). **3a** was stable under general conditions in which common acetal survive [with piperidine (base), Bu₄NF (F⁻), and LiAlH₄ (hydride)]. Interestingly, **3a** withstood mild acidic conditions (Table 3, Runs 1–4). Deprotection reaction of **3a** was successfully proceeded under stronger acidic conditions (2 M HCl aq. in 1,4-dioxane at 80 °C) or oxidation conditions with aqueous ceric ammonium nitrate (CAN) to afford corresponding ketone **2a** in high yields (Runs 5 and 6).^[13]

Table 2. PdBr₂-acetalization of various aldehydes and ketones with diazophenanthrenequinone **1**.^[a]

[a] Reaction conditions: **2/1/ PdBr₂** = 1/1.5/0.1. Benzene (0.2 M for **2**). Reflux for 1.5 h.

Table 3. Examination of the stability of acetal **3a** under various conditions.^[a]

Run	Conditions	Temp	Time	Yield (%)	
				2a	3a
1	<i>p</i> -TsOH (20 mol%), THF/H ₂ O = 1:1	r.t.	5	0	88
2	2 M HCl aq./1,4-dioxane = 1:1	r.t.	5	0	90
3	10% H ₂ SO ₄ aq./THF = 1:1	r.t.	5	0	90
4	CH ₃ CO ₂ H	r.t.	5	0	92
5	2 M HCl aq./1,4-dioxane = 1/1	80 °C	8	82	6
6 ^[b]	CAN (2.0 eq.), CH ₃ CN/H ₂ O = 1/1	0 °C	5	84	0

[a] Reaction was carried out for 5 h. [b] CAN: (NH₄)₂[Ce(NO₃)₆].

In Table 4, the results of the deprotection reaction of various 9,10-phenanthrenedioxy acetals **3** are shown. Since several carbonyl compounds **2** were over oxidized with the aqueous CAN solution, the yields were low (Runs 3 and 6).


Table 4. Deprotection of acetal **3**.^[a]

Run	Acetal 3	Product 2	Yield (%) [Time (h)]	
			Method A	Method B
1	3a	2a	82 [8]	84 [5]
2	3c	2c	80 [1]	80 [2.5]
3	3e	2e	82 [12]	40 [9] ^[b]
4	3i	2i	90 [5] ^[c]	100 [5]
5	3n	2n	24 [15] ^[d]	78 [1.5] ^[b]
6	3q	2q	80 [3]	38 [6] ^[b]

[a] Reaction conditions. Method A: In 1,4-dioxane/2 M HCl=1/1 at 80 °C. Method B: CAN (2 equiv.) in CH₃CN/H₂O=1/2 at 0 °C. [b] In CHCl₃/H₂O=1/2. [c] In CHCl₃/2 M HCl=1/1. [d] **3n** was recovered in 63% yield.

COMMUNICATION

In Scheme 4, demonstration of synthetic utility of the 9,10-phenanthrenedioxy acetal are shown. When 2 M HCl aqueous solution was treated to a mixture of 9,10-phenanthrenedioxy acetal **3d** and ethyleneglycol acetal **7** at room temperature in THF, ethyleneglycol acetal **7** was selectively deprotected and 9,10-phenanthrenedioxy acetal was isolated in 84% yield. **3h** possesses two kinds of acid sensitive protective groups such as *tert*-butyldimethylsilyl group and 9,10-phenanthrenedioxy acetal, these protective groups are selectively removed by treatment using tetrabutylammonium fluoride and CAN to afford phenol **8** and ketone **2h**, respectively.



Scheme 4. Demonstration of selective deprotection.

In conclusion, we developed the PdBr₂-catalyzed acetal formation reaction with diazophenanthrenequinone under non-acidic conditions. The acetal that has a phenanthrene skeleton has good stability against mild acidic reaction conditions and was transformed to the corresponding carbonyl compound under strong acidic or oxidation conditions using aqueous CAN.


Acknowledgements

This work was supported by a Grant in Aid for KAKENHI (17K19125).

Keywords: acetal • diazo compounds • palladium • protective group

- [1] a) M. B. Smith, J. March, *March's Advanced Organic Chemistry 6th ed.* John Wiley & Sons, Inc., New Jersey, **2007**, 587-655. b) G. Brahmachari, *RCS Adv.* **2016**, 6, 64676. c) C. J. Li, *Chem. Rev.* **1993**, 93, 2023.
- [2] P. G. M. Wuts, T. W. Greene, *Greene's Protective groups in organic synthesis 5th ed.* John Wiley & Sons, Inc., New Jersey, **2014**.
- [3] a) M. K. Basu, S. Samajdar, F. F. Becker, B. K. Banik, *Synlett* **2002**, 319. b) B. Karimi, B. Golshani, *Synthesis* **2002**, 784.
- [4] a) J. Grabowski, J. M. Granda, J. Jurczak, *Org. Biomol. Chem.* **2018**, 16, 3114. b) J. Deutsch, A. Martin, H. Lieske, *J. Catal.* **2007**, 245, 428. c) M. Barbasiewicz, M. Maukosza, *Org. Lett.* **2006**, 8, 3745.
- [5] a) C. Tortoreto, T. A. L. Egger, L. Guenee, J. Lacour, *Org. Lett.* **2016**, 18, 240. b) M. E. Alonso, M. C. Garcia, A. W. Chitty, *J. Org. Chem.* **1985**, 50, 3445. c) Z. Yang, C. Lei, X. Zhao, R. Liu, H. Wei, Y. Ma, S. Meng, Q. Cao, J. Wei, X. Wang, *ChemistrySelect* **2017**, 2, 9377. d) S. K. De, R. A. Gibbs, *Tetrahedron Lett.* **2004**, 45, 8141. e) A. Clerici, N. Pastori, O. Porta, *Tetrahedron* **1998**, 54, 15679.
- [6] a) A. Dhakshinamoorthy, M. Alvaro, M. Puche, V. Fornes, H. Garcia, *ChemCatChem* **2012**, 4, 2026. b) M. X. Tan, L. Gu, N. Li, J. Y. Ying, Y. Zhang, *Green Chem.* **2013**, 15, 1127.
- [7] a) For review, see: D. I. A. Othman, M. Kitamura, *Heterocycles* **2016**, 92, 1761.
- [8] Recent publication of our group, see: a) M. Kitamura, R. Sakata, T. Okauchi, *Tetrahedron Lett.* **2011**, 52, 1931; b) M. Kitamura, M. Kisanuki, R. Sakata, T. Okauchi, *Chem. Lett.* **2011**, 40, 1129; c) M. Kitamura, M. Kisanuki, T. Okauchi, *Eur. J. Org. Chem.* **2012**, 905; d) M. Kitamura, K. Kubo, S. Yoshinaga, *Tetrahedron Lett.* **2014**, 55, 1653; e) M. Kitamura, M. Kisanuki, K. Kanemura, T. Okauchi, *Org. Lett.* **2014**, 16, 1554; f) M. Kitamura, S. Takahashi, T. Okauchi, *J. Org. Chem.* **2015**, 80, 8406; g) M. Kitamura, K. Otsuka, S. Takahashi, T. Okauchi, *Tetrahedron Lett.* **2017**, 58, 3508; h) D. I. A. Othman, K. Otsuka, S. Takahashi, K. B. Selim, M. A. El-Sayed, A. S. Tantawy, T. Okauchi, M. Kitamura, *Synlett* **2018**, 29, 457; i) S. Takahashi, H. Shimooka, T. Okauchi, M. Kitamura, *Chem. Lett.* **2019**, 48, 28. Recent reaction of diazophenanthrenequinone by other groups, see: j) K. Bahadur, S. Magar, Y. R. Lee, *Org. Lett.* **2013**, 15, 4288; k) E. K. R. Baral, Y. R. Lee, S. H. Kim, *Adv. Synth. Catal.* **2015**, 357, 2883; l) E. K. R. Baral, Y. R. Lee, S. H. Kim, Y. J. Wee, *Synthesis* **2016**, 48, 579; m) K. B. Somai Magar, T. N. J. I. Edison, Y. R. Lee, *Eur. J. Org. Chem.* **2017**, 7046; n) B. Ghosh, A. Biswas, S. Chakraborty, R. Samanta, *Chem. - Asian J.* **2018**, 13, 2388; o) H.-X. Wang, Q. Wan, K. Wu, K.-H. Low, C. Yang, C.-Y. Zhou, J.-S. Huang, C.-M. Che, *J. Am. Chem. Soc.* **2019**, 141, 9027; p) H.-X. Wang, Y. Richard, Q. Wan, C.-Y. Zhou, C.-M. Che, *Angew. Chem., Int. Ed.* **2020**, 59, 1845.
- [9] For reviews, see: a) M. P. Doyle, *Chem. Rev.* **1986**, 86, 919; b) A. Padwa, K. E. Krumpe, *Tetrahedron* **1992**, 48, 5385; c) T. Ye, M. A. McKervey, *Chem. Rev.* **1994**, 94, 1091; d) A. Padwa, D. J. Austin, *Angew. Chem., Int. Ed. Engl.* **1994**, 33, 1797; e) M. P. Doyle, T. Ye, M. A. McKervey, in *Modern catalytic methods for organic synthesis with diazo compounds*, John Wiley & Sons: New York, **1998**; f) H. M. L. Davies, R. E. J. Beckwith, *Chem. Rev.* **2003**, 103, 2861; g) Z. Zhang, J. Wang, *Tetrahedron* **2008**, 64, 6577; h) M. P. Doyle, R. Duffy, M. Ratnikov, L. Zhou, *Chem. Rev.* **2010**, 110, 704-724. i) A. Ford, H. Miel, A. Ring, C. N. Slattery, A. R. Maguire, M. A. McKervey, *Chem. Rev.* **2015**, 115, 9981.
- [10] For a review, see: A. Padwa, S. F. Hornbuckle, *Chem. Rev.* **1991**, 91, 263.
- [11] a) M. Kitamura, N. Tashiro, R. Sakata, T. Okauchi, *Synlett* **2010**, 2503; b) M. Kitamura, R. Sakata, N. Tashiro, A. Ikegami, T. Okauchi, *Bull. Chem. Soc. Jpn.* **2015**, 88, 824.
- [12] J. E. Baldwin, *J. Chem. Soc. Chem. Commun.* **1976**, 734.
- [13] 9,10-Phenanthrenequinone was formed as by-product in the deprotection reaction.

Entry for the Table of Contents



A new acetalization method of ketones and aldehydes under non-acidic conditions was developed using diazophenanthrenequinone and PdBr₂. The formed acetals that have a phenanthrene skeleton withstand under mild acidic conditions. Removal of acetals was successfully proceeded under strong acidic or oxidation conditions using aqueous ceric ammonium nitrate (CAN).

Key topic

Acetal Formation